

Large N reduction for Chern-Simons theory on S^3

Goro Ishiki^{1,*}, Shinji Shimasaki^{2,†} and Asato Tsuchiya^{3‡}

¹ *High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305-0801, Japan*

² *Department of Physics, Kyoto University, Kyoto 606-8502, Japan*

³ *Department of Physics, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan*

(Dated: August, 2009; preprint: KEK-TH-1326, KUNS-2226)

We study a matrix model which is obtained by dimensional reduction of Chern-Simon theory on S^3 to zero dimension. We find that expanded around a particular background consisting of multiple fuzzy spheres, it reproduces the original theory on S^3 in the planar limit. This is viewed as a new type of the large N reduction generalized to curved space.

PACS numbers: 11.25.Tq; 11.15.Tk; 11.10.Kk

Introduction.— Emergent space(-time) is often seen in recent developments in string theory. In particular, it is characteristic of the matrix models [1] which are dimensionally reduced models of ten-dimensional super Yang-Mills theory (SYM) and proposed as a non-perturbative formulation of superstring or M-theory. While it is verified that these models reproduce gravity around flat space-time, it is necessary to elucidate how curved space-time is included in these models [2]. The large N reduction [4] developed decades ago is the first example that realizes emergent space-time in matrix model. Indeed, it asserts that the planar limit of gauge theory on flat space-time is equivalent to the planar limit of a matrix model (a reduced model) that is obtained by dimensional reduction to lower dimensions. It is, therefore, worthwhile to study generalization of the large N reduction to curved space-time. The large N reduction can also give a non-perturbative formulation of planar gauge theory as an alternative to lattice gauge theory. It is well-known that in order to overcome the problem of flat directions (or the $U(1)^D$ symmetry breaking) in the large N reduction, one needs the prescription [5], which is unfortunately not compatible with supersymmetry in gauge theory.

The authors of Ref. [6] proposed a non-perturbative formulation of planar $\mathcal{N} = 4$ SYM on $R \times S^3$ equivalent to that on R^4 at a conformal point, using the plane wave matrix model (PWMM) [7] (See also [8, 9]). PWMM is obtained by dimensional reduction of $\mathcal{N} = 4$ SYM on $R \times S^3$ to R [10], and in the formulation the S^3 is realized as a non-trivial fiber bundle over S^2 by expanding PWMM around a particular background which consists of multiple fuzzy spheres. Thus the formulation is viewed as a new type of the large N reduction generalized to curved space. The formulation overcomes the aforementioned problem in the large N reduction for supersymmetric gauge theory thanks to massiveness and supersymmetry of PWMM [11]. By putting the formulation on a computer in terms of the method [13], one should be able to study the strongly coupled regime of $\mathcal{N} = 4$ SYM and therefore to perform new non-trivial tests for the AdS/CFT duality [14].

A reasonable argument supporting the validity of the

proposal was given in [6], and the proposal has already passed some non-trivial tests at weak coupling [6, 15]. Application of the same type of the large N reduction to various gauge theories was also discussed in [16]. However, the proposal has not been proved completely yet, in particular at strong coupling. In this letter, we prove that the same type of the large N reduction does work even at strong coupling for another gauge theory, Chern-Simons (CS) theory on S^3 , which has been exactly solved [17]. Our study is an extension of the classical analysis done in [18] to the quantum level and provides the first proved example of the new type of the large N reduction. Our formalism gives a new quantum mechanical definition of CS theory on S^3 , which is an interesting topological field theory associated with the knot theory. CS theory on S^3 is also interpreted as open topological A strings on T^*S^3 . This aspect should enable us to gain from our findings insights into the study of formulating superstring non-perturbatively in terms of matrix model.

CS theory on S^3 .— Let us review some exact results for $U(N)$ CS theory on S^3 , whose action is given by

$$S_{CS} = \frac{k}{4\pi} \int_{S^3} \text{Tr} \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right). \quad (1)$$

The partition function of the theory, $\mathcal{Z}_{CS}$, defines a topological invariant of the manifold S^3 . Given an oriented knot $\mathcal{K}$ in S^3 , one can consider the Wilson loop in an irreducible representation R of $U(N)$

$$W_R(\mathcal{K}) = \text{Tr}_R \left(P \exp \oint_{\mathcal{K}} A \right). \quad (2)$$

The expectation value of the Wilson loop, $\langle W_R(\mathcal{K}) \rangle_{CS}$, defines a topological invariant of $\mathcal{K}$. In this letter, we mainly consider the Wilson loop for an unknot in the fundamental representation, denoted by $W_{\square}(\text{unknot})$, where $\square$ stands for the fundamental representation. It turns out that our formalism adopts a non-canonical framing corresponding to the one labeled by $m = n = 1$ in [19].

In our framing, the above quantities are given by [17, 19, 20]

$$\mathcal{Z}_{CS} = e^{-\frac{1}{12} g_s k (N^2 - 1)} \left(\frac{g_s}{2\pi i} \right)^{N/2} \prod_{\alpha > 0} 2 \sinh \frac{g_s \alpha \cdot \rho}{2} \quad (3)$$

$$= \int \prod_{i=1}^N \frac{d\beta_i}{2\pi} \prod_{i < j} \sinh^2 \frac{\beta_i - \beta_j}{2} e^{-\frac{1}{2g_s} \sum_i \beta_i^2}. \quad (4)$$

$$\langle W_{\square}(\text{unknot}) \rangle_{CS} = e^{\frac{g_s}{2}(N - \frac{1}{N})} \frac{\sinh \frac{g_s N}{2}}{\sinh \frac{g_s}{2}} = \sum_{i=1}^N \langle e^{\beta_i} \rangle_{CSM}, \quad (5)$$

where ρ is the Weyl vector of $SU(N)$, $\alpha > 0$ are positive roots, and we have introduced $g_s = 2\pi i/(k+N)$, which is identified with the string coupling in topological strings. We have ignored a g_s -independent factor in rewriting (3) to (4). $\langle \cdots \rangle_{CSM}$ denotes the expectation value with the weight given by the integrand in (4).

Dimensional reduction.— In order to dimensionally reduce CS theory on S^3 [18, 21], we regard S^3 as the $SU(2)$ group manifold, which has the isometry $SO(4) = SU(2) \times SU(2)$ corresponding to the left and right translations. We set the radius of S^3 to $2/\mu$, and define the right-invariant 1-forms E^i ($i = 1, 2, 3$), which satisfy the Maurer-Cartan equation $dE^i - \frac{\mu}{2} \epsilon_{ijk} E^j \wedge E^k = 0$. We define the Killing vector $\mathcal{L}_i$ that is dual to E^i and generates $SU(2)$ of the left translation. We can also regard S^3 as an $S^1(U(1))$ bundle over $S^2 = SU(2)/U(1)$. The Kaluza-Klein (KK) momenta q in the fiber direction S^1 take integers and half-integers. When $\mathcal{L}_i$ act on a KK mode with the momentum q , it takes the form of the angular momentum operators $L_i^{(q)}$ in the presence of a monopole with magnetic charge q [22]. Namely, the KK momenta in the fiber direction are identified with the monopole charges on S^2 .

Expanding the gauge field in (1) as $A = i\mu X_i E^i$, we rewrite RHS of (1) as

$$-\frac{2k}{\pi} \int d\Omega_3 \text{Tr} \left(i\epsilon^{ijk} \left(X_i \mathcal{L}_j X_k + \frac{2}{3} X_i X_j X_k \right) + X_i^2 \right), \quad (6)$$

where $d\Omega_3$ is the volume element of unit 3-sphere, and we have used the Maurer-Cartan equation. By only keeping the $q = 0$ modes in (6), namely replacing $\mathcal{L}_i$ and $d\Omega_3$ with $L_i^{(0)}$ and $d\Omega_2$, respectively, we can dimensionally reduce the theory onto S^2 . The resultant theory is Yang-Mills theory (YM) on S^2 . To see this, we decompose X_i into the radial component χ and the components tangential to S^2 , a_θ and a_φ , and rewrite the theory on S^2 as $\int d\Omega_2 \text{Tr} (\chi \epsilon^{\mu\nu} f_{\mu\nu} - \chi^2)$, where $f_{\mu\nu}$ ($\mu, \nu = \theta, \varphi$) is the field strength for a_μ . Integrating χ out indeed yields YM on S^2 . Finally, dropping all the derivatives in (6), namely dimensionally reducing the theory onto a point gives rise to a three-matrix model:

$$S_m = -\frac{1}{g_m^2} \text{Tr} \left(X_i^2 + \frac{i}{3} \epsilon^{ijk} X_i [X_j, X_k] \right). \quad (7)$$

YM on S^2 from the matrix model.— The matrix model (7) with the matrix size $M \times M$ possesses the following classical solutions,

$$\hat{X}_i = \bigoplus_s L_i^{[j_s]} \otimes \mathbf{1}_{N_s}, \quad (8)$$

where $L_i^{[j]}$ are the spin j representation of the $SU(2)$ generators, the relation $\sum_s (2j_s + 1)N_s = M$ is satisfied, and $j_s \neq j_t$ for $s \neq t$. The label s runs over some integers.

The (s, t) block of the fluctuation around (8) is expanded in terms of the fuzzy spherical harmonics [8, 9, 23] which is a $(2j_s + 1) \times (2j_t + 1)$ matrix. We put $2j_s + 1 = \Omega + n_s$ with Ω and n_s being integers and take the limit in which

$$\Omega \rightarrow \infty, \quad g_m^2/\Omega = -g_{YM}^2 A/(8\pi^2) \text{ fixed} \quad (9)$$

with $g_{YM}^2 = -\mu^2/(2k)$. $A = 4\pi/\mu^2$ is the area of the S^2 , and g_{YM} turns out to be the coupling constant of YM on S^2 . In this limit, the above fuzzy spherical harmonics coincides classically with the monopole harmonics [24] of the monopole charge $j_s - j_t$ under the identification $L_i^{[j_s]} \bullet \cdots \bullet L_i^{[j_t]} \leftrightarrow L_i^{(j_s - j_t)} \bullet$ and $\frac{1}{\Omega} \text{Tr} \leftrightarrow \frac{1}{4\pi} \int d\Omega_2$. It was indeed shown in [18] that the theory around (8) is classically equivalent to the theory around the following classical solution of $U(K)$ YM on S^2 ,

$$\hat{X}_i = \text{diag}(\cdots, \underbrace{L_i^{(q_s)} - L_i^{(0)}, \dots, L_i^{(q_s)} - L_i^{(0)}}_{N_s}, \cdots) \quad (10)$$

expressed in terms of X_i , where $q_s = n_s/2$, and the relation $\sum_s N_s = K$ is satisfied.

The above relationship is extended to the quantum level [21]. The partition function of the matrix model (7) is decomposed into sectors classified by the representation of $SU(2)$ as in (8), and in the limit (9) the sectors consisting of K blocks reduces to

$$\sum_{\{n_s, N_s\}} \int \prod_s \prod_{i=1}^{N_s} dy_{si} \prod_{s \leq t} \prod_{i=1}^{N_s} \prod_{j=1}^{N_t} \{ (y_{si} - y_{tj})^2 - (n_s - n_t)^2 \} \\ \times e^{-\frac{2\pi^2}{g_{YM}^2 A} \sum_s \sum_{i=1}^{N_s} (y_{si}^2 + n_s^2)} \quad (11)$$

up to an irrelevant constant with an analytic continuation $g_{YM}^2 \rightarrow -ig_{YM}^2$. y_{si} come from the eigenvalues of X_3 . This exactly agrees with the partition function of $U(K)$ 2d YM on S^2 [25, 26], and the $\{n_s, N_s\}$ sector in the summation in (11) is the contribution around the classical background (10).

CS theory on S^3 from the matrix model.— CS theory on S^3 is obtained by summing all the KK modes in YM on S^2 , where the KK momenta are identified with the monopole charges on S^2 . In the limit (9), the (s, t) block of the fluctuation around (8) of the matrix model (7) behaves as the one around (10) of YM on S^2 which feels the monopole charge $j_s - j_t$. Therefore we are led to the following statement: if one chooses in (8)

$$-\Lambda/2 \leq s \leq \Lambda/2, \quad 2j_s + 1 = \Omega + s, \quad N_s = N, \quad (12)$$

with Λ being a positive even integer and takes the limit in which

$$\Omega \rightarrow \infty, \quad \Lambda \rightarrow \infty, \quad \Omega - \frac{\Lambda}{2} \rightarrow \infty, \quad N \rightarrow \infty,$$

$$g_m^2 N / \Omega = N / (4\pi(k + N)) \text{ fixed}, \quad (13)$$

the theory around (12) of the matrix model (7) is equivalent to $U(N)$ CS theory on S^3 in the planar limit. The naive relation between the coupling constants is $g_m^2 N / \Omega = N / (4\pi k)$, but we will see below that this naive one is renormalized so that the last equation in (13) is valid. The particular background (12) and the limit (13) is the same as the ones adopted in [6] in realizing $\mathcal{N} = 4$ SYM on $R \times S^3$ in PWMM. This extraction of the theory around (12) from the whole theory of the matrix model should be allowed in the $N \rightarrow \infty$ limit as in the case of PWMM. The above equivalence was classically shown in [18] by further imposing the orbifolding condition, which needs infinitely large matrix size and is not allowed quantum mechanically. Here we do not need to impose it because of the $N \rightarrow \infty$ limit. Ω and Λ are viewed as the ultraviolet cutoffs for the angular momenta, which remarkably respect the gauge symmetry.

More precisely, the equivalence states that

$$\mathcal{F}_m / (N^2(\Lambda + 1)) = \mathcal{F}_{CS} / N^2 \quad (14)$$

in the limit (13), where $\mathcal{F}_m$ is the free energy of the theory around the background (12) of the matrix model (7), and $\mathcal{F}_{CS} = \ln \mathcal{Z}_{CS}$. Furthermore, given a knot $\mathcal{K}$ in S^3 , we can parametrize it by $z^M(\sigma)$ ($\sigma \in [0, 1]$) and introduce a Wilson loop in the matrix model,

$$\hat{W}(\mathcal{K}) = \frac{1}{M} \text{Tr} \left[P \exp \left(i\mu \int_0^1 X_i E_M^i(z(\sigma)) \frac{dz^M(\sigma)}{d\sigma} d\sigma \right) \right].$$

It was shown classically in [18] that the above operator around (12) in the limit (13) corresponds to $W_\square(\mathcal{K})$ in (2). We naturally expect this correspondence to hold also at the quantum level:

$$\langle \hat{W}(\mathcal{K}) \rangle_m = \langle W_\square(\mathcal{K}) \rangle_{CS} / N, \quad (15)$$

where $\langle \cdots \rangle_m$ denotes the expectation value around the background (12) of the matrix model in the limit (13). Here we concentrate on an unknot whose path is given by a great circle on S^3 . In this case, the relation (15) reduces to

$$\langle \text{Tr} (e^{4\pi i X_3}) \rangle_m / M = \langle W_\square(\text{unknot}) \rangle_{CS} / N. \quad (16)$$

Proof of the equivalence. — To prove the equivalence, we show the equalities (14) and (16) explicitly. For that purpose, we may use the statistical model whose partition function is defined by (11), by taking the $\Omega \rightarrow \infty$ limit first. We extract the contribution of $-\Lambda/2 \leq s \leq \Lambda/2$, $n_s = s$, $N_s = N$ from the sum in (11). Then, we see that the model is interpreted as a multi-matrix model of $N \times N$ hermitian matrices with double trace interactions in which y_{si} is a constant times the i -th eigenvalue of the s -th matrix ϕ_s :

$$\mathcal{Z} = \int \prod_s d\phi_s \exp \left(-\frac{N}{2} \sum_s \text{tr} \phi_s^2 - \tilde{V} \right), \quad (17)$$

where

$$\tilde{V} = \sum_{s \neq t} \sum_{k=1}^{\infty} \sum_{m=0}^{2k} \frac{1}{2k(s-t)^2} \left(\frac{-\lambda}{4\pi^2} \right)^k \binom{2k}{m} \times \text{tr} \phi_s^m \text{tr} (-\phi_t)^{2k-m}$$

with $\lambda = g_s N = 2\pi i N / (k + N)$. (13) tells us to take the limit in which $\Lambda \rightarrow \infty$, $N \rightarrow \infty$ with λ fixed. Recalling the computation in obtaining (11) from the matrix model (7) in [21], we find that LHS of (16) is rewritten as $\langle \sum_s \text{tr} e^{\sqrt{\lambda} \phi_s} / (N(\Lambda + 1)) \rangle$, where $\langle \cdots \rangle$ denotes the expectation value in (17). In the same way, by using the formula $\prod_{n=1}^{\infty} (1 + x^2/n^2) = \sinh(\pi x) / (\pi x)$, we also see that the statistical model defined by (4) is interpreted as a one-matrix model of an $N \times N$ hermitian matrix ϕ with double-trace interaction terms [27]:

$$\mathcal{Z}_M = \int d\phi \exp \left(-\frac{N}{2} \text{tr} \phi^2 - V \right), \quad (18)$$

where

$$V = \sum_{k=1}^{\infty} \sum_{m=0}^{2k} \frac{\zeta(2k)}{k} \left(\frac{-\lambda}{4\pi^2} \right)^k \binom{2k}{m} \text{tr} \phi^m \text{tr} (-\phi)^{2k-m}.$$

Noting (5), we finally see that (14) and (16) reduce to

$$\mathcal{F} / (N^2(\Lambda + 1)) = \mathcal{F}_M / N^2, \quad (19)$$

$$\frac{1}{N(\Lambda + 1)} \sum_s \langle \text{tr} e^{\sqrt{\lambda} \phi_s} \rangle = \frac{1}{N} \langle \text{tr} e^{\sqrt{\lambda} \phi} \rangle_M, \quad (20)$$

respectively, where $\mathcal{F} = \ln \mathcal{Z}$, $\mathcal{F}_M = \ln \mathcal{Z}_M$, and $\langle \cdots \rangle_M$ denotes the expectation value in (18).

To show (19) and (20), we perform the perturbative expansion in λ in both the matrix models using the standard double line notation. We represent the vertices by connecting the double traces in terms of dashed lines as in Fig. 1. There is one-to-one correspondence between the diagrams of the two matrix models. As in the standard perturbation theory, the free energy and the expectation value of the Wilson loop are computed by summing all the connected diagrams, where the diagrams that have pieces connected in terms of the dashed lines are regarded as connected. It is easy to see that the leading contribution in the $1/N$ expansion is given by the diagrams that are planar in the ordinary sense and in addition divided into two parts by cutting any dashed line. The diagrams that do not satisfy these two conditions give the subleading contribution in the $1/N$ expansion. Namely, only such ‘tree’ planar diagrams survive in the $N \rightarrow \infty$ limit. Fig. 1 shows examples of the diagrams for the free energy: the left one is ‘tree’ planar and gives the leading contribution, while the right one gives the subleading contribution. For the left diagram in Fig. 1, we can prove an equality

$$\frac{1}{2^4(\Lambda + 1)} \sum_{s,t,u,v,w} \frac{1}{(s-t)^8(t-u)^{10}(u-v)^4(u-w)^4}$$




FIG. 1: Diagrams for the free energy: A ‘tree’ planar diagram giving the leading contribution in the $1/N$ expansion (left) and a diagram giving the subleading contribution (right). Each number represents the degree of the vertex.




FIG. 2: A generic ‘tree’ planar diagram for the free energy. Each blob represents a planar diagram.

$$= \zeta(8)\zeta(10)\zeta(4)\zeta(4),$$

in the $\Lambda \rightarrow \infty$ limit, so that we find that the diagram takes the same value in the two matrix models. Such an equality does not hold for the right diagram in Fig. 1, so that the diagram takes different values in the two matrix models. For a generic ‘tree’ planar diagram shown in Fig. 2 possessing n dashed lines, we can prove the following equality in the $\Lambda \rightarrow \infty$ limit [28]:

$$\frac{1}{2^n(\Lambda + 1)} \sum_{\{s_a\}} \frac{1}{(s_1 - s_2)^{2k_1} \cdots (s_n - s_{n+1})^{2k_n}} = \prod_{a=1}^n \zeta(2k_a),$$

which implies that the diagram takes the same value in the two matrix models. Thus we have shown that (19) and (20) hold to all order in λ in the limit in which $\Lambda \rightarrow \infty$ and $N \rightarrow \infty$.

Summary.— In this letter, we showed that expanded around the background (12) in the limit (13), the matrix model is equivalent to planar $U(N)$ CS theory on S^3 . We can also show that expanded around different backgrounds, the matrix model reproduces planar CS theory on S^3/Z_q , which we will describe in [28]. Our formalism does not only give a new regularization method of planar CS theory on S^3 , but also serves as the first proved example of the new type of the large N reduction.

Acknowledgments.— We would like to thank K. Ohta for his collaboration in the early stage of this work and for many discussions. The work of S. S. is supported by JSPS. The work of A. T. is supported by Grant-in-Aid for Scientific Research (19540294) from JSPS.

- [†] Electronic address: shinji@gauge.scphys.kyoto-u.ac.jp
- [‡] Electronic address: satsuch@ipc.shizuoka.ac.jp
- [1] T. Banks, W. Fischler, S. H. Shenker and L. Susskind, Phys. Rev. D **55**, 5112 (1997); N. Ishibashi, H. Kawai, Y. Kitazawa and A. Tsuchiya, Nucl. Phys. B **498**, 467 (1997); R. Dijkgraaf, E. P. Verlinde and H. L. Verlinde, Nucl. Phys. B **500**, 43 (1997).
- [2] For an interesting approach to this subject, see Ref. [3].
- [3] M. Hanada, H. Kawai and Y. Kimura, Prog. Theor. Phys. **114**, 1295 (2006).
- [4] T. Eguchi and H. Kawai, Phys. Rev. Lett. **48**, 1063 (1982).
- [5] G. Bhanot et al., Phys. Lett. B **113**, 47 (1982); D. J. Gross and Y. Kitazawa, Nucl. Phys. B **206**, 440 (1982); A. Gonzalez-Arroyo and M. Okawa, Phys. Rev. D **27**, 2397 (1983).
- [6] T. Ishii, G. Ishiki, S. Shimasaki and A. Tsuchiya, Phys. Rev. D **78**, 106001 (2008).
- [7] D. E. Berenstein, J. M. Maldacena and H. S. Nastase, JHEP **0204**, 013 (2002).
- [8] G. Ishiki, S. Shimasaki, Y. Takayama and A. Tsuchiya, JHEP **0611**, 089 (2006).
- [9] T. Ishii, G. Ishiki, S. Shimasaki and A. Tsuchiya, Phys. Rev. D **77**, 126015 (2008).
- [10] N. Kim et al., Nucl. Phys. B **671**, 359 (2003).
- [11] For recent developments in alternative lattice approach to supersymmetric gauge theory, see Refs. [12].
- [12] I. Kanamori, Phys. Rev. D **79**, 115015 (2009); S. Catterall et al., arXiv:0903.4881 [hep-lat]; D. Kadoh et al., Nucl. Phys. B **820**, 99 (2009); A. D’Adda et al., arXiv:0907.4137 [hep-th]; M. Hanada and I. Kanamori, arXiv:0907.4966 [hep-lat].
- [13] K. N. Anagnostopoulos, M. Hanada, J. Nishimura and S. Takeuchi, Phys. Rev. Lett. **100**, 021601 (2008); S. Catterall and T. Wiseman, Phys. Rev. D **78**, 041502 (2008).
- [14] J. M. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998).
- [15] G. Ishiki, S. W. Kim, J. Nishimura and A. Tsuchiya, Phys. Rev. Lett. **102**, 111601 (2009), arXiv:0907.1488 [hep-th]; Y. Kitazawa and K. Matsumoto, arXiv:0811.0529 [hep-th].
- [16] M. Hanada, L. Mannelli and Y. Matsuo, arXiv:0905.2995 [hep-th], arXiv:0907.4937 [hep-th].
- [17] E. Witten, Commun. Math. Phys. **121**, 351 (1989).
- [18] T. Ishii, G. Ishiki, K. Ohta, S. Shimasaki and A. Tsuchiya, Prog. Theor. Phys. **119**, 863 (2008).
- [19] M. Blau and G. Thompson, JHEP **0605**, 003 (2006).
- [20] M. Marino, Commun. Math. Phys. **253**, 25 (2004), Rev. Mod. Phys. **77**, 675 (2005), arXiv:hep-th/0410165; M. Tierz, Mod. Phys. Lett. A **19**, 1365 (2004).
- [21] G. Ishiki, K. Ohta, S. Shimasaki and A. Tsuchiya, Phys. Lett. B **672**, 289 (2009).
- [22] For explicit forms of $\mathcal{L}_i$ and $L_i^{(q)}$, see Refs. [6, 9, 18, 21].
- [23] H. Grosse et al., Commun. Math. Phys. **178**, 507 (1996); K. Dasgupta et al., JHEP **0205**, 056 (2002).
- [24] T. T. Wu and C. N. Yang, Nucl. Phys. B **107**, 365 (1976).
- [25] A. A. Migdal, Sov. Phys. JETP **42**, 413 (1975); J. A. Minahan and A. P. Polychronakos, Nucl. Phys. B **422**, 172 (1994).
- [26] Indeed, only the $SU(K)$ part agrees [21]. However, the difference does not affect the following analysis.
- [27] M. Aganagic et al., JHEP **0402**, 010 (2004).
- [28] G. Ishiki, S. Shimasaki and A. Tsuchiya, in preparation.

* Electronic address: ishiki@post.kek.jp